

급성 심근 경색증환자에서의 심근 관류의 평가

— 심근 조영심초음파를 이용한 연구 —

임세중¹ · 하중원¹ · 최동훈¹ · 고충원¹ · 이동일¹ · 편옥범¹
전국진¹ · 안신기¹ · 이문형¹ · 장양수¹ · 이종두² · 정남식¹

= Abstract =

Assessment of Myocardial Perfusion in Patients with Acute Myocardial Infarction

- Myocardial Contrast Echocardiography Using Intravenous Infusion of Microbubbles -

Se-Joong Rim, MD¹, Jong-Won Ha, MD¹, Dong-Hoon Choi, MD¹,
Choong-Won Goh, MD¹, Dong-Il Lee, MD¹, Wook-Bum Pyun, MD¹,
Kook Jin Chun, MD¹, Shinki Ahn, MD¹, Moon-Hyoung Lee, MD¹,
Yangsoo Jang, MD¹, Jong Doo Lee, MD² and Namsik Chung, MD¹

¹Cardiology Division, Department of Internal Medicine, ²Nuclear Medicine,
Yonsei University College of Medicine, Seoul, Korea

Background : Unlike ^{99m}Tc-Sestamibi, microbubbles used during myocardial contrast echocardiography (MCE) exist only in the vascular space. Therefore, there may be a difference in the pattern of myocardial perfusion between MCE and ^{99m}Tc-Sestamibi Single-Photon Emission Computed Tomography (SPECT) in acute myocardial infarction (AMI). **Objectives** : The purpose of this study was to assess myocardial perfusion using MCE with intravenous infusion of perfluorocarbon-exposed sonicated dextrose albumin microbubbles (IV MCE), and to compare it with SPECT and MCE with intracoronary injection of sonicated Hexabrix (IC MCE). **Methods** : Seventeen patients with AMI (male 13, age 59.5 ± 8.8 years, anterior MI 10) underwent IV MCE at 8.1 ± 3.7 days after onset. SPECT and IC MCE were also performed at 1.2 ± 1.0 days and 2.0 ± 1.5 days from IV MCE respectively. Any revascularization procedures were not performed between three studies. Perfusion defect by three methods was scored semiquantitatively as 1 : normal perfusion, 0.5 : moderate defect, and 0 : severe defect at 16 segments of the left ventricle. **Results** : 1) Perfusion defect in infarction territory was detected in 15 patients with SPECT, 12 patients with IV MCE and 11 patients with IC MCE. 2) Concordance of perfusion score at each segment was 93% between IV MCE and IC MCE, 65% between IV

: 2001 4 25
: 2001 5 20
: , 120 - 752 134
: (02) 361 - 7071 · : (02) 393 - 2041 E-mail : sejoong@yumc.yonsei.ac.kr

KEY WORDS : Myocardial contrast echocardiography · Acute myocardial infarction.

IV MCE

가 peak videointen-
sity

(microbubble) (imaging technique) . (IC MCE)가

IV

, IV MCE

(IV MCE) 가 .

IV MCE

가
가 IC MCE 가
가 가

17

	IV	(creat-
MCE	^{99m} Tc - Sestamibi inine phosphokinase)	13
Single - Photon Emission Computed Tomography (SPECT)	가 , ± 8.8 . 10 , 7	59.5
IV MCE	/	

IV MCE가 SPECT MCE가 가 .

1~

가

4

IV MCE perfluorocarbon - exposed sonicated dextrose albumin microbubble(PESDA)²⁻⁴⁾ . Apical 4,3,2 chamber view baseline PESDA intermittent Harmonic imaging pulsing interval 1 : 1, 1 : 4, 1 : 7 가 (1) moderate defect(0.5), severe defect(0) (Fig. 1). off - line vide- view continuous imaging videotape . PESDA 8 mL perfluor-

opropane (MW 188 g/mol), 12 mL 5% dextrose water, 4 mL 5% human albumin . 80 sonication (550 W : 25%). PESDA 0.05 mL/kg 100 mL 0.5~ 15 mL/min . IC MCE Hexabrix[®] sonication 2 mL . apical 4,3,2 chamber view continuous imaging videotape 1 ,




Fig. 1. Apical 2 chamber view during intermittent harmonic imaging. With increasing pulsing interval from 1 : 1 to 1 : 4 and 1 : 7, perfusion defect in apical area (arrow) became partially enhanced.




Fig. 2. Comparison of intravenous myocardial contrast echocardiography (IV MCE), ^{99m}Tc-Sestamibi SPECT (SPECT), and intracoronary myocardial contrast echocardiography (IC MCE) using left coronary artery(LCA) injection. Apical 4 chamber view imagings of IV and IC MCE and corresponding horizontal long axis view of SPECT showed perfusion defects in apical area.

가 0.5 , 가 0 SPECT ^{99m}Tc -Sestamibi short, horizontal long, vertical long axis uptake 100%(1),




Fig. 3. Discrepancy of perfusion defect observed between MCE and SPECT. A : Apical 4 chamber view of IV and IC MCE (left coronary artery injection) and corresponding horizontal long axis view of SPECT in one patient with anterior wall myocardial infarction. B : Apical 2 chamber view of IV and IC MCE (right coronary artery injection) and corresponding vertical long axis view of SPECT in one patient with inferior wall myocardial infarction. C : Apical 4 chamber view of IV and IC MCE (left coronary artery injection) and corresponding horizontal long axis view of SPECT in one patient with lateral wall myocardial infarction (See text for details).

Table 1. Detection of perfusion defect within infarction territory using three different methods

Patient No.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17
SPECT	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	-	-
IC MCE	+	+	+	+	+	+	+	+	+	+	+	-	-	-	-	-	-
IV MCE	+	+	+	+	+	+	+	+	+	+	+	+	-	-	-	-	-

+ : perfusion defect, - : normal perfusion

25~75%(0.5), 25% (0)

3가

16 2

-

결 과

1) IV MCE 8.1 ± 3.7 (4~20)

. IV MCE SPECT

1.2 ± 1.0 , IV MCE IC MCE

2.0 ± 1.5 .

2) IV MCE

$9.7 \pm 2.4(4 \sim 13)$

mL/minute.

3) IV MCE, IC MCE, SPECT

가 Fig. 2

4) MCE SPECT

Fig. 3. Fig. 3A

SPECT, IV MCE IC MCE

. Fig. 3B

SPECT

IV IC MCE

가 Fig. 3C

MCE SPECT

5) 17

SPECT 15

MCE 11, IV MCE 12, IC MCE (Table 1).

6)

MCE 93% IV MCE IC MCE (Table 2).

IV MCE (0 0.5)가

Table 2. Observed agreement between IV MCE and IC MCE

		IC MCE			
		Severe defect	Mild defect	No defect	Total
IV MCE	Severe defect	12	5	4	21
	Mild defect	7	8	2	17
	No defect	0	0	234	234
	Total	19	13	240	272

Concordance 93%

Table 3. Observed agreement between IV MCE and SPECT

		SPECT			
		Severe defect	Mild defect	No defect	Total
IV MCE	Severe defect	11	5	5	21
	Mild defect	7	8	2	17
	No defect	22	55	157	234
	Total	40	68	164	272

Concordance 65%

Table 4. Observed agreement between IC MCE and SPECT

		SPECT			
		Severe defect	Mild defect	No defect	Total
IC MCE	Severe defect	11	6	2	19
	Mild defect	4	6	3	13
	No defect	25	56	159	240
	Total	40	68	164	272

Concordance 64%

32 IC MCE 가

7) IV MCE SPECT 65%, IC MCE

SPECT 64% MCE SPECT

(Table 3 and 4). SPECT

108 31 IV MCE

고 안

가⁹⁾가
MCE SPECT 가
가 (TIMI SPECT imaging partial volume
grade 3) 가 effect 가¹⁰⁾
“ no - reflow ” SPECT
“ no - reflow ”가⁵⁾
가 videointensity uptake
가⁶⁾⁷⁾ “ no -
reflow ”
IC MCE
IV MCE
가⁸⁾
가 IC MCE IV MCE,
SPECT가
no - reflow가 MCE
가 가
no - reflow 가

요 약

no - 연구목적 :
reflow no - reflow가^{99m}Tc -
MCE SPECT Sestamibi
가 가
Perfluorocarbon - exposed sonicated dextrose
albumin (IV
MCE) 가
SPECT가^{99m}Tc - Sestamibi SPECT sonicated Hexabrix®
(IC MCE)
MCE 가 SPECT
gold standard
대상 및 방법 :
가 17 (13 , 59.5 ±
8.8) 4~20(8.1 ± 3.7) IV
MCE 4 SPECT
MCE 1 (1.2 ± 1.0) IC MCE(2.0 ± 1.5)
16
MCE 1 : , 0.5 : ,
0 :

결 과 :

1) 17
SPECT 15 , IV MCE 12 , IC MCE 11

2) IV MCE IC MCE 93%, IV MCE
SPECT 65%, IC MCE SPECT 64%

3) IC MCE 32
IV MCE
, SPECT
108 31 IV MCE

결 론 :

가 IV MCE IC MCE
SPECT
가

중심 단어 :

2001

REFERENCES

- 1) Marwick TH, Brunken R, Meland N, et al. Accuracy and feasibility of contrast echocardiography for detection of perfusion defects in routine practice: comparison with wall motion and MIBI-SPECT. *J Am Coll Cardiol* 1998; 32:1260-9.
- 2) Porter TR, Xie F. Transient myocardial contrast following initial exposure to diagnostic ultrasound pressures with minute doses of intravenously injected microbubbles: demonstration and potential mechanisms. *Circulation* 1995; 92:2391-5.

- 3) Porter TR, Xie F, Kricsfeld D, Armbruster RW. Improved myocardial contrast with second harmonic transient ultrasound contrast response imaging in humans using intravenous perfluorocarbon-exposed sonicated dextrose albumin. *J Am Coll Cardiol* 1996;27:1497-501.
- 4) Porter TR, Li S, Kricsfeld D, Armbruster RW. Detection of myocardial perfusion in multiple echocardiographic windows with one intravenous injection of microbubbles using transient response second harmonic imaging. *J Am Coll Cardiol* 1997;29:791-3.
- 5) Ito H, Okamura A, Iwakura K, Masuyama T, Hori M, Takiuchi S, Negoro S, Nakatsuchi Y, Taniyama Y, Higashino Y, Fujii K, Minamino T. Myocardial perfusion patterns related to thrombolysis in myocardial infarction perfusion grades after coronary angioplasty in patients with acute anterior wall myocardial infarction. *Circulation* 1996;93:1993-9.
- 6) Ito H, Maruyama A, Iwakura K, Takiuchi S, Masuyama T, Hori M, Higashino Y, Fujii K, Minamino T. Clinical implications of the 'no reflow' phenomenon. A predictor of complications and left ventricular remodeling in reperfused anterior wall myocardial infarction. *Circulation* 1996;93: 223-8.
- 7) Ragosta M, Camarano G, Kaul S, Powers ER, Sarembock IJ, Gimble LW. Microvascular integrity indicates myocellular viability in patients with recent myocardial infarction. *Circulation* 1994;89:2562-9.
- 8) Thomas R Porter, Shouping Li, Rebecca Oster. Ubeydullah Deligonul The clinical implications of no reflow demonstrated with intravenous perfluorocarbon containing microbubbles following restoration of thrombolysis in myocardial infarction (TIMI) 3 flow in patients with acute myocardial infarction. *Am J Cardiol* 1997;82 (10): 1173-7.
- 9) Ito H, Iwakura K, Oh H, Masuyama T, Hori M, Higashino Y, Fujii K, Minamino T. Temporal changes in myocardial perfusion patterns in patients with reperfused anterior wall myocardial infarction. Their relation to myocardial viability. *Circulation* 1995;91:656-62.
- 10) Pretorius PH, King MA, Pan TS, de Vries DJ, Glick SJ, Byrne CL. Reducing the influence of the partial volume effect on SPECT activity quantitation with 3D modelling of spatial resolution in iterative reconstruction. *Physics in Medicine & Biology* 1998;43:407-20.